

Spin and quadrupole correlations in the insulating nematic phase of spin-1 bosons in optical lattices

Guang-Ming Zhang¹ and Lu Yu²

¹*Department of Physics and Center for Advanced Study, Tsinghua University, Beijing 100084, China;*

²*Institute of Theoretical Physics and Interdisciplinary Center of Theoretical Studies, CAS, Beijing 100080, China.*

(Dated: February 2, 2008)

We consider the effective model of $H = -J_1 \sum_{\langle i,j \rangle} \mathbf{S}_i \cdot \mathbf{S}_j - J_2 \sum_{\langle i,j \rangle} (\mathbf{S}_i \cdot \mathbf{S}_j)^2$, describing the Mott insulating phase with odd number of spin-1 bosons in optical lattices ($J_2 > J_1 > 0$). In terms of an SU(3) boson representation, a valence bond mean field theory is developed. In 1D, a *first-order* quantum phase transition from a spin singlet to a spin nematic phase with *gapful* excitations is identified at $J_1/J_2 = 0.19833$, while on a 2D square lattice a spin nematic ordered phase with gapless excitations prevails. In both 1D and 2D cases, we predict that the spin structure factor displays dominant antiferromagnetic fluctuations, while the quadrupole structure factor exhibits strong ferroquadrupolar correlations.

PACS numbers: 03.75.Mn, 32.80.Pj, 75.10.Jm

Degenerate alkali atoms are considered weakly interacting boson gas due to the smallness of the scattering length compared with the inter-particle separation. However, the situation changes dramatically when an optical potential created by standing laser beams confines particles in valleys of the periodic potential and strongly enhances local interactions. Recently, the nontrivial Mott insulating state of bosonic atoms in optical lattices has been demonstrated experimentally [1]. In addition to solid state systems, spinor atoms in optical lattices provide a novel realization of quantum magnetic systems with the possibility to tune various parameters of the effective models in the absence of disorder [2, 3, 4].

Alkali atoms have a nuclear spin 3/2. Lower energy hyperfine manifold has three magnetic sublevels and a total moment $S = 1$. In order to observe the quantum spin phenomena experimentally, one has to consider a system with *small* number of particles and *strong* interactions. In the insulating state, atoms are localized, and fluctuations in the particle number on each site are suppressed. Virtual tunnelling of atoms between neighboring sites induces effective spin interactions, leading to novel quantum magnetic phases [5, 6, 7, 8].

The boson-Hubbard model is used to describe the low-energy physics of spin-1 bosonic atoms in an optical lattice [9],

$$H = -t \sum_{\langle i,j \rangle, m} (a_{i,m}^\dagger a_{j,m} + h.c.) + \frac{U_0}{2} \sum_i n_i(n_i - 1) + \frac{U_2}{2} \sum_i (\mathbf{S}_i^2 - 2n_i) - \mu \sum_i n_i, \quad (1)$$

where $n_i = \sum_m a_{i,m}^\dagger a_{i,m}$ and $S_i^\alpha = \sum_{m,n} a_{i,m}^\dagger T_{m,n}^\alpha a_{i,n}$ denote the number of atoms and the spin operator on the i site ($m = -1, 0, 1$), respectively, while T^α denote the matrices for the spin-1 particles.

In the $t = 0$ limit with *odd* number of bosons per site, the bosonic symmetry of the wave function requires

that each site has a localized spin with *odd* quantum numbers. The interactions of Eq.(1) are minimized when the localized spins take the smallest possible value $S = 1$. Away from this limit, the second order perturbation theory in t leads to an effective spin superexchange model [6, 7, 10]

$$H = -J_1 \sum_{\langle i,j \rangle} \mathbf{S}_i \cdot \mathbf{S}_j - J_2 \sum_{\langle i,j \rangle} (\mathbf{S}_i \cdot \mathbf{S}_j)^2, \quad (2)$$

where the coupling parameters are confined to $0 < J_1/J_2 < 1$. For a two-spin problem, the first term favors a polarized configuration, while the second prefers a spin singlet form. Thus, the effective model acts as a spin-1 Heisenberg *antiferromagnet* with ferromagnetic frustrations.

It has been an outstanding issue to study the ground states of this effective model [6, 7, 8, 11, 12]. More exotic ground states may be realized in 1D [8, 13, 14], but there is no consensus yet. In this paper, using an SU(3) boson representation, we develop a valence bond mean field (MF) theory. In 1D, a *first-order* quantum phase transition is found from a spin singlet ($J_1/J_2 < 0.19833$) to a (short-range ordered) spin nematic phase ($J_1/J_2 > 0.19833$) with gapful excitations, while on a 2D square lattice an *ordered* spin nematic phase is always obtained. Moreover, the spin and quadrupole correlation spectra in both disordered and ordered nematic phases are calculated explicitly, to be checked by future experiments.

To describe the spin-1 operators, an SU(3) bosonic representation is defined by generators $F_m^n(i) = a_{i,m}^\dagger a_{i,n}$, where indices m and n specify the spin projection with values $-1, 0, 1$. The commutation relation is satisfied

$$[F_m^n(i), F_\mu^\nu(j)] = \delta_{i,j} [\delta_{n,\mu} F_m^\nu(i) - \delta_{m,\nu} F_\mu^n(i)], \quad (3)$$

which forms an SU(3) Lie algebra. The corresponding spin operators are expressed as $S_i^+ = \sqrt{2}(a_{i,0}^\dagger a_{i,-1} + a_{i,1}^\dagger a_{i,0})$, $S_i^- = (S_i^+)^\dagger$, $S_i^z = (a_{i,1}^\dagger a_{i,1} - a_{i,-1}^\dagger a_{i,-1})$. With

this boson representation, we can verify $[S_i^+, S_j^-] = 2S_i^z \delta_{i,j}$, $[S_i^z, S_j^\pm] = \pm S_i^\pm \delta_{i,j}$. In order to fix $S_i^2 = S(S+1) = 2$, a local constraint $\sum_m a_{i,m}^\dagger a_{i,m} = 1$ has to be imposed.

To make the spin representation symmetric, linear combinations are introduced: $b_{i,1} = \frac{1}{\sqrt{2}}(a_{i,-1} - a_{i,1})$, $b_{i,2} = \frac{-i}{\sqrt{2}}(a_{i,-1} + a_{i,1})$, and $b_{i,3} = a_{i,0}$, and the spin (dipolar) operators are written in antisymmetric form

$$\begin{aligned} S_i^x &= -i(b_{i,2}^\dagger b_{i,3} - b_{i,3}^\dagger b_{i,2}), \\ S_i^y &= -i(b_{i,3}^\dagger b_{i,1} - b_{i,1}^\dagger b_{i,3}), \\ S_i^z &= -i(b_{i,1}^\dagger b_{i,2} - b_{i,2}^\dagger b_{i,1}), \end{aligned} \quad (4)$$

and the local constraint holds as $\sum_\alpha b_{i,\alpha}^\dagger b_{i,\alpha} = 1$. Actually for spin-1 bosonic atoms, *quadrupole* operators can be defined by

$$\begin{aligned} Q_i^{(0)} &= 3(S_i^z)^2 - 2 = b_{i,1}^\dagger b_{i,1} + b_{i,2}^\dagger b_{i,2} - 2b_{i,3}^\dagger b_{i,3}, \\ Q_i^{(2)} &= (S_i^x)^2 - (S_i^y)^2 = -(b_{i,1}^\dagger b_{i,1} - b_{i,2}^\dagger b_{i,2}), \\ Q_i^{xy} &= S_i^x S_i^y + S_i^y S_i^x = -(b_{i,1}^\dagger b_{i,2} + b_{i,2}^\dagger b_{i,1}), \\ Q_i^{xz} &= S_i^x S_i^z + S_i^z S_i^x = -(b_{i,1}^\dagger b_{i,3} + b_{i,3}^\dagger b_{i,1}), \\ Q_i^{yz} &= S_i^y S_i^z + S_i^z S_i^y = -(b_{i,2}^\dagger b_{i,3} + b_{i,3}^\dagger b_{i,2}). \end{aligned} \quad (5)$$

Three dipole and five quadrupole operators form generators of the SU(3) Lie group, as in the Gell-Mann matrix representation. Then, the effective spin model is expressed as:

$$\begin{aligned} H &= - \sum_{\langle i,j \rangle} \sum_{\alpha,\beta} \left[J_1 b_{i,\alpha}^\dagger b_{i,\beta} b_{j,\beta}^\dagger b_{j,\alpha} \right. \\ &\quad \left. + (J_2 - J_1) b_{i,\alpha}^\dagger b_{i,\beta} b_{j,\alpha}^\dagger b_{j,\beta} \right]. \end{aligned} \quad (6)$$

In the limit $J_1 = J_2$, the model is reduced to an SU(3) symmetric *ferromagnetic* superexchange model, invariant under the uniform SU(3) transformation, for which the spin ferromagnetic and spin nematic long-range order can coexist in higher dimensions [15]. On the other hand, in the limit $J_1 = 0$, the model is reduced to an SU(3) symmetric valence-bond *antiferromagnetic* model, invariant under the staggered conjugate transformations of the two sublattices, leading to a spin dimerization in 1D [16, 17].

It is known that an SU(2) Schwinger boson MF theory can describe rather well the spin correlations for spin-1/2 Heisenberg antiferromagnets [18]. To develop a similar valence bond MF theory, we introduce symmetric pairing parameters

$$\begin{aligned} \Delta_{\alpha,\alpha} &= -\langle b_{i,\alpha}^\dagger b_{j,\alpha}^\dagger \rangle, \\ \Delta_{1,2} &= -\langle b_{i,1}^\dagger b_{j,2}^\dagger + b_{i,2}^\dagger b_{j,1}^\dagger \rangle, \\ \Delta_{1,3} &= -\langle b_{i,1}^\dagger b_{j,3}^\dagger + b_{i,3}^\dagger b_{j,1}^\dagger \rangle, \\ \Delta_{2,3} &= -\langle b_{i,2}^\dagger b_{j,3}^\dagger + b_{i,3}^\dagger b_{j,2}^\dagger \rangle. \end{aligned} \quad (7)$$

To preserve the SU(2) spin rotational symmetry of the model, we have to assume $\Delta_{1,1} = \Delta_{2,2} = \Delta_{3,3} \equiv \Delta_a$ and $\Delta_{1,2} = \Delta_{1,3} = \Delta_{2,3} \equiv \Delta_b$. By introducing a Nambu spinor $\Psi_{\mathbf{k}} = (b_{\mathbf{k},1}^\dagger, b_{\mathbf{k},2}^\dagger, b_{\mathbf{k},3}^\dagger, b_{-\mathbf{k},1}, b_{-\mathbf{k},2}, b_{-\mathbf{k},3})$, we can rewrite the MF Hamiltonian in a matrix form

$$\begin{aligned} H_{mf} &= \frac{1}{2} \sum_{\mathbf{k}} \Psi_{\mathbf{k}}^\dagger \mathbf{H}_{mf}(\mathbf{k}) \Psi_{\mathbf{k}} \\ &\quad + \frac{3}{2} [(3J_2 - J_1) \Delta_a^2 + J_1 \Delta_b^2] Nz - \frac{5}{2} \lambda N, \end{aligned} \quad (8)$$

where the local constraint has been implemented by a Lagrangian multiplier λ , N the total number of lattice sites, and z the number of the nearest neighbor sites. The MF Hamiltonian matrix is given by

$$\mathbf{H}_{mf}(\mathbf{k}) = \lambda + 2\Delta_a(\mathbf{k})\sigma_x \otimes I + \Delta_b(\mathbf{k})\sigma_x \otimes M, \quad (9)$$

where $M = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}$, I is the 3×3 unit matrix, σ_α ($\alpha = x, y, z$) are Pauli matrices, $\Delta_a(\mathbf{k}) = \frac{z}{2} \Delta_a (3J_2 - J_1) \gamma_{\mathbf{k}}$, $\Delta_b(\mathbf{k}) = z \Delta_b J_1 \gamma_{\mathbf{k}}$, $\gamma_{\mathbf{k}} = \frac{1}{d} \sum_{\delta} \cos \mathbf{k} \cdot \delta$, and δ the nearest neighbor vector. The corresponding Matsubara Green function (GF) is thus deduced to

$$\mathbf{G}^{-1}(\mathbf{k}, i\omega_n) = i\omega_n \sigma_z \otimes I - \mathbf{H}_{mf}(\mathbf{k}), \quad (10)$$

where ω_n is the bosonic Matsubara frequency. The poles of the GF matrix give rise to the quasiparticle spectra: $\epsilon_1(\mathbf{k}) = \sqrt{\lambda^2 - 4[\Delta_a(\mathbf{k}) + \Delta_b(\mathbf{k})]^2}$ and $\epsilon_2(\mathbf{k}) = \sqrt{\lambda^2 - [2\Delta_a(\mathbf{k}) - \Delta_b(\mathbf{k})]^2}$, where the lower band $\epsilon_1(\mathbf{k})$ is singly occupied while the higher band $\epsilon_2(\mathbf{k})$ is doubly degenerate.

From the free energy of the system, the saddle point equations at $T = 0\text{K}$ are derived as

$$\begin{aligned} \frac{1}{N} \sum_{\mathbf{k}} \left[\frac{\lambda}{2\epsilon_1(\mathbf{k})} + \frac{\lambda}{\epsilon_2(\mathbf{k})} \right] &= \frac{5}{2}, \\ \frac{1}{N} \sum_{\mathbf{k}} \frac{\gamma_{\mathbf{k}}^2}{\epsilon_1(\mathbf{k})} &= \frac{2(\Delta_a + \Delta_b)}{z[(3J_2 - J_1)\Delta_a + 2J_1\Delta_b]}, \\ \frac{1}{N} \sum_{\mathbf{k}} \frac{\gamma_{\mathbf{k}}^2}{\epsilon_2(\mathbf{k})} &= \frac{2\Delta_a - \Delta_b}{z[(3J_2 - J_1)\Delta_a - J_1\Delta_b]}, \end{aligned} \quad (11)$$

while the ground state energy is $E_g = -\frac{3}{2}zN[(3J_2 - J_1)\Delta_a^2 + J_1\Delta_b^2]$. From the GF matrix, we can also derive the double-time GFs of the boson operators

$$\begin{aligned} \langle\langle b_{\mathbf{k},2} | b_{\mathbf{k},1}^\dagger \rangle\rangle &= \langle\langle b_{\mathbf{k},3} | b_{\mathbf{k},1}^\dagger \rangle\rangle = \langle\langle b_{\mathbf{k},3} | b_{\mathbf{k},2}^\dagger \rangle\rangle \\ &= \frac{-(i\omega_n + \lambda)[4\Delta_a(\mathbf{k})\Delta_b(\mathbf{k}) + \Delta_b^2(\mathbf{k})]}{[\omega_n^2 + \epsilon_1^2(\mathbf{k})][\omega_n^2 + \epsilon_2^2(\mathbf{k})]}, \end{aligned} \quad (12)$$

and $\langle\langle b_{\mathbf{k},1} | b_{\mathbf{k},1}^\dagger \rangle\rangle = \langle\langle b_{\mathbf{k},2} | b_{\mathbf{k},2}^\dagger \rangle\rangle = \langle\langle b_{\mathbf{k},3} | b_{\mathbf{k},3}^\dagger \rangle\rangle$. Using the spectral representation, we can calculate the expectation

values at $T = 0\text{K}$,

$$\begin{aligned} \langle b_{i,1}^\dagger b_{i,2} \rangle &= \langle b_{i,1}^\dagger b_{i,3} \rangle = \langle b_{i,2}^\dagger b_{i,3} \rangle \\ &= \frac{1}{6N} \sum_{\mathbf{k}} \left[\frac{\lambda}{\epsilon_1(\mathbf{k})} - \frac{\lambda}{\epsilon_2(\mathbf{k})} \right], \end{aligned} \quad (13)$$

and $\langle b_{i,1}^\dagger b_{i,1} \rangle = \langle b_{i,2}^\dagger b_{i,2} \rangle = \langle b_{i,3}^\dagger b_{i,3} \rangle = 1/3$. Therefore, we find $\langle S_i^\alpha \rangle = 0$ and $\langle (S_i^\alpha)^2 \rangle = 2/3$ ($\alpha = x, y, z$), exhibiting that both time reversal and spin rotational symmetries are well preserved in the present MF theory. Then the spin nematic state with a quadrupole moment $\langle Q_i^{xy} \rangle = \langle Q_i^{xz} \rangle = \langle Q_i^{yz} \rangle \equiv -Q$ is a possible ground state.

To evaluate the spin spatial correlations in the Mott insulating phase, the dynamic correlation functions should be calculated. By expressing the spin operators in terms of the Nambu spinor, the spin correlations are given by

$$\begin{aligned} &\chi^{\alpha,\beta}(\mathbf{q}, i\omega_m) \\ &= \frac{1}{4\beta N} \sum_{\mathbf{k}, \omega_n} \text{Tr} [\Gamma_\alpha \mathbf{G}(\mathbf{k}, i\omega_n) \Gamma_\beta \mathbf{G}(\mathbf{k} + \mathbf{q}, i\omega_n + i\omega_m)], \end{aligned}$$

where $\Gamma_x, \Gamma_y, \Gamma_z$ denote the corresponding 6×6 matrices of S_i^x, S_i^y, S_i^z , respectively, and both ω_n and ω_m are bosonic Matsubara frequencies. Inserting the GF matrix and tracing over matrices, we find

$$\chi^{x,x}(\mathbf{q}, i\omega_m) = \chi^{y,y}(\mathbf{q}, i\omega_m) = \chi^{z,z}(\mathbf{q}, i\omega_m), \quad (14)$$

displaying an $SU(2)$ spin rotational symmetry. The corresponding imaginary part $\chi''(\mathbf{q}, \omega)$ can also be obtained through performing the summation over the Matsubara frequency and analytic continuation [19]. According to the fluctuation-dissipation theorem, the static spin structure factor is thus obtained $S_D(\mathbf{q}) = \frac{-1}{\pi} \int_{-\infty}^{+\infty} d\omega [1 + n_B(\omega)] \chi''(\mathbf{q}, \omega)$. Furthermore, when the correlation function of the quadrupole operator is defined by $S_Q(\mathbf{r} - \mathbf{r}') = \langle [3(S_r^z)^2 - 2][3(S_{r'}^z)^2 - 2] \rangle$, the static quadrupole structure factor $S_Q(\mathbf{q})$ can be evaluated as well [19].

In 1D, $z = 2$ and $\gamma_{\mathbf{k}} = \cos k$. For a given value of J_1/J_2 , the saddle point equations are solved numerically, results being displayed in Fig.1. For $0 \leq J_1/J_2 < 0.19833$, we find $\Delta_b = 0$, $\Delta_a = 0.5077J_2$, the quasiparticle band $\epsilon_1(k)$ and $\epsilon_2(k)$ are degenerate, so the ground state is a spin singlet with an energy gap. From the point of view of spin correlations, such a spin singlet phase is similar to the spin dimerized state in the limit $J_1 = 0$ [17]. For $J_1/J_2 > 0.19833$, both Δ_b and Δ_a are finite, and two energy gaps in the quasiparticle bands are found $\Delta_{s,1} = \sqrt{\lambda^2 - 4[(3J_2 - J_1)\Delta_a + 2J_1\Delta_b]^2}$ and $\Delta_{s,2} = \sqrt{\lambda^2 - 4[(3J_2 - J_1)\Delta_a - J_1\Delta_b]^2}$ at momenta $k = 0, \pm\pi$. Moreover, the quadrupole moment Q jumps from 0 to 0.16897 at the critical coupling $J_1/J_2 = 0.19833$, and then the system is in a short-range ordered spin singlet phase with a *finite* quadrupole moment. Thus, there is a *first-order* quantum phase transition from the spin singlet to spin nematic phase with gapful excitations.




FIG. 1: Saddle point solution in 1D is displayed: the valence bond parameters (a), the ground state energy (b), two energy gaps of quasiparticle bands (c), and the quadrupole moment (d) as functions of J_1/J_2 . J_2 is taken as the energy unit.

If we rewrite $J_1 = -\cos \theta$ and $J_2 = -\sin \theta$, the critical coupling corresponds to $\theta_c = -0.5633\pi$. In fact, a *disordered* spin nematic (non-dimerized) phase *with breaking the $SU(2)$ spin rotational symmetry* was first suggested around $\theta \approx -3\pi/4$ by Chubukov [13]. However, subsequent numerical work [14] did not support this proposal and it was then believed that the dimerized phase prevails in the whole regime $-3\pi/4 \leq \theta \leq -\pi/2$, i.e., $0 < J_1/J_2 < 1$. On the other hand, recent quantum Monte Carlo [20], density matrix renormalization group calculations [21], and quantum field theory approach [22] indicate that the dimerized phase may end at $\theta \sim -0.67\pi$, casting doubts on the earlier anticipations.

To put our MF results on a solid ground, the static spin structure factor $S_D(q)$ and quadrupole structure factor $S_Q(q)$ are calculated at $T = 0\text{K}$ and displayed in Fig.2. We find that $S_D(q)$ exhibits a broad peak around the antiferromagnetic wave vector $q = \pi$ in both spin singlet and gapped spin nematic phase. However, the quadrupole structure factor $S_Q(q)$ shows a broad peak around the ferromagnetic wave vector $q = 0$ in the spin singlet phase, while $S_Q(q)$ exhibits a *sharp* resonance at $q = 0$ in the spin nematic phase, indicating strong *ferro-quadrupolar* spatial correlations.

On a 2D square lattice, we have $z = 4$ and $\gamma_{\mathbf{k}} = (\cos k_x + \cos k_y)/2$. At $T = 0\text{K}$, the conversion from summation over momentum to integral will be invalid as $\lambda \rightarrow 4[(3J_2 - J_1)\Delta_a + 2J_1\Delta_b]$, and then the quasiparticle band $\epsilon_1(\mathbf{k})$ becomes gapless and linear near $\mathbf{k}^* = (0, 0)$ and (π, π) . The Bose-Einstein condensation thus occurs, while a finite energy gap still exists in the band $\epsilon_2(\mathbf{k})$. By separating the divergent term from the summation, we can introduce a superfluid density ρ_s and then solve the saddle point equations. The results are




FIG. 2: Spin structure factor (a) and quadrupole structure factors (b) in 1D for spin singlet phase (solid line $J_1/J_2 = 0.1$) and spin nematic phase (dashed line $J_1/J_2 = 0.5$).




FIG. 3: Saddle point solution on a 2D square lattice is shown: the valence bond parameters (a), the superfluid density and quadrupole moment (b) as functions of J_1/J_2 .

shown in Fig.3, where a finite quadrupole moment Q is always obtained and it increases as the coupling J_1/J_2 grows. The ground state in the regime $0 \leq J_1/J_2 \leq 1$ is thus a long-range *ordered* spin nematic phase.

It is also more important to examine the spin and quadrupole spatial correlations in the ordered nematic phase. The calculated static spin structure factor $S_D(q_x, q_y)$ is shown in Fig.4a, where a strong broad peak is displayed at $\mathbf{q} = (\pi, \pi)$, indicating dominant antiferromagnetic spin correlations, and weaker broad peaks at $\mathbf{q} = (0, \pi)$ and $\mathbf{q} = (\pi, 0)$, corresponding to collinear

spin-dimer correlations. As a comparison, the calculated static quadrupole structure factor $S_Q(q_x, q_y)$ is displayed in Fig.4b, where a δ -like resonance appears at $\mathbf{q} = (0, 0)$, implying strong *ferroquadrupolar* long-range correlations. These are unique features of an ordered spin nematic phase, which can be directly probed by polarized inelastic light scattering experimentally.

To summarize, the SU(3) boson representation is used to develop an efficient valence bond MF theory for the Mott insulation phase with odd number of spin-1 bosons in optical lattices. A *first-order* quantum phase transition from a spin singlet to a nematic phase with gapful excitations is predicted for 1D, while on a 2D square lattice a spin nematic *ordered* phase is shown to always prevail. Both predictions are further supported by an explicit calculation of the spin and quadrupole correlation functions.

G.-M. Zhang is grateful to H. -F. Lü for his help in numerical calculations, and acknowledges the support of NSF-China (No.10125418 and 10474051) and the Special




FIG. 4: Spin structure factor (a) and quadrupole structure factor (b) for the 2D spin nematic ordered phase with $J_1/J_2 = 0.3$.

Fund for Major State Basic Research Projects of China (No.G200067107). L. Yu also acknowledges the support of NSF-China.

-
- [1] M. Greiner *et al.*, Nature **415**, 39 (2002).
 - [2] D. Jaksch *et al.*, Phys. Rev. Lett. **81**, 3108 (1998).
 - [3] L.-M. Duan, E. Demler, and M. D. Lukin, Phys. Rev. Lett. **91**, 090402 (2003).
 - [4] J. J. García-Ripoll, M. A. Martin-Delgado, and J. I. Cirac, Phys. Rev. Lett. **93**, 250405 (2004).
 - [5] E. Demler and F. Zhou, Phys. Rev. Lett. **88**, 163001 (2002).
 - [6] M. Snoek and F. Zhou, Phys. Rev. B **69**, 094410 (2004).
 - [7] A. Imambekov, M. Lukin, and E. Demler, Phys. Rev. A **68**, 063602 (2003); Phys. Rev. Lett. **93**, 120405 (2004).
 - [8] S. K. Yip, Phys. Rev. Lett. **90**, 250402 (2003).
 - [9] T. L. Ho, Phys. Rev. Lett. **81**, 742 (1998); T. Ohmi and K. Machida, J. Phys. Soc. Jpn. **67**, 1822 (1998).
 - [10] S. Tsuchiya, S. Kurihara, and T. Kimura, Phys. Rev. A **70**, 043628 (2004).
 - [11] H. H. Chen and P. M. Levy, Phys. Rev. B **7**, 4267 (1973); N. Papanicolaou, Nucl. Phys. B **305**, 367 (1988).
 - [12] K. Harada and N. Kawashima, Phys. Rev. B **65**, 052403 (2002).
 - [13] A. V. Chubukov, Phys. Rev. B **43**, 3337 (1991).
 - [14] G. Fáth and J. Sólyom, Phys. Rev. B **51**, 3620 (1995).
 - [15] C. D. Batista, G. Ortiz, and J. E. Gubernatis, Phys. Rev. B **65**, 180402 (2002).
 - [16] I. Affleck, Nucl. Phys. B **265**, 409 (1986).
 - [17] M. N. Barber and M. T. Batchelor, Phys. Rev. B **40**, 4621 (1989); A. Klumper, Europhys. Lett. **9**, 815 (1989).
 - [18] D. P. Arovas and A. Auerbach, Phys. Rev. B **38**, 316 (1988); Phys. Rev. Lett. **61**, 617 (1988).
 - [19] G. -M. Zhang *et al.*, in preparation.
 - [20] N. Kawashima, Prog. Theor. Phys. Suppl. **145**, 138 (2002).
 - [21] A. Läuchli, G. Schmid, and S. Trebst, cond-mat/0311082.
 - [22] B. A. Ivanov and A. K. Kolezhuk, Phys. Rev. B **68**,

052401 (2003).